J. of Ramanujan Society of Math. and Math. Sc. Vol.4, No.2 (2015), pp. 71-76

Higher order derivative on meromorphic functions in terms of subordination

ISSN: 2319-1023

Sh. Najafzadeh

Department of Mathematics, Pavame Noor University, Iran

Email addresses: najafzadeh1234@yahoo.ie

Dedicated to Prof. Hari M. Srivastava on his 75th birth anniversary

Abstract: This paper investigates sharp coefficient bounds, integral representation, extreme point and operator properties of a certain class associated with functions which are meromorphic in the punctured unit disk.

keywords and phrases: Meromorphic function, Coefficient estimate, Extreme point, Integral representation, Integral operator.

2000 Mathematics Subject Classification: 30C45, 30C50.

1. Introduction

Let \sum denote the class of meromorphic functions of the form

$$f(z) = \frac{1}{z} + \sum_{n=3}^{\infty} a_n z^n \quad (a_n \ge 0, \quad n \in \mathbb{N}),$$
 (1.1)

which are analytic and univalent in the punctured unit disk $\Delta^* = \{z: 0 < |z| < 1\}$.

For $f(z) \in \Sigma$, Ghanim and Darus [3] were defined a linear operator I^k (k = 0, 1, 2, ...) as follows:

$$I^{0}f(z) = f(z)$$

$$I^{k}f(z) = z(I^{k-1}f(z))' + \frac{2}{z} = \frac{1}{z} + \sum_{n=3}^{+\infty} n^{k} a_{n} z^{n}.$$
(1.2)

For A = B + (C - B)(1 - D), $-1 \le B < C \le 1$ and $0 \le D < 1$, we let $\sum_{A,B}^{K}$ consists of function $f \in \sum$ satisfying the condition

$$-\frac{zF^{(4)}(z)}{F'''(z)} < 4\frac{1+Az}{1+Bz},\tag{1.3}$$

where $F(z) = I^k f(z)$ is defined by (1.2).

For other subclass of meromorphic univalent functions, we can see the recent works of many authors in [1] and [2].