

**A NOTE ON CERTAIN IDENTITIES INVOLVING GENERAL
DOUBLE AND TRIPLE HYPERGEOMETRIC FUNCTIONS
OF EXTON AND SRIVASTAVA**

M. I. Qureshi¹, K. A. Quraishi^{2,*}, Mohd. Sadiq Khan³ and A. Arora⁴

¹Department of Applied Sciences and Humanities, Faculty of Engineering and
Technology, Jamia Millia Islamia (A Central University),
New Delhi-110025 (India).

²Mathematics Section, Mewat Engineering College (Waqf),
Palla, Nuh, Mewat, Haryana-122107 (India).

³Department of Mathematics, Shibli National College,
Azamgarh, Uttar Pradesh-276001 (India).

⁴Department of Applied Sciences and Humanities, Accurate Institute of
Engineering and Technology, Knowledge Park-III, Greater Noida, Gautam
Buddha Nagar, U.P.-201306 (India).

E-mails:

miqureshi_delhi@yahoo.co.in; kaleemspn@yahoo.co.in; msadiqkhan.snc@gmail.com

*Corresponding Author

Abstract: The main object of the present note is to obtain the generalizations and unifications of identities associated with hypergeometric function of one variable ${}_A F_B$ and Appell's hypergeometric function of two variables F_1, F_2, F_3, F_4 due to MacRobert and Sharma respectively, in the form of identities associated with multiple hypergeometric functions \mathcal{H}, G of Exton and $F^{(3)}$ of Srivastava.

Keywords and Phrases: Pochhammer's symbol; Multiplication formulae, Double and triple hypergeometric functions; Series iteration technique.

2010 Mathematics Subject Classification: 33C05, 33C20, 33C65, 33C70.

1. Introduction

In 1967, a unification and generalization of Lauricella's fourteen complete triple hypergeometric functions of second order F_1, F_2, \dots, F_{14} [8, pp. 113-114] including Saran's ten triple hypergeometric functions $F_E, F_F, F_G, F_K, F_M, F_N, F_P, F_R, F_S,$