

CERTAIN RESULTS ON CONTINUED FRACTIONS

Satya Prakash Singh and Ashish Pratap Singh

Department of Mathematics,
T. D. P. G. College, Jaunpur,
Jaunpur - 222002, Uttar Pradesh, INDIA

E-mail : snsp39@gmail.com, ashishsingh1671@gmail.com

(Received: May 31, 2023 Accepted: Jun. 18, 2023 Published: Jun. 30, 2023)

Abstract: In this paper certain results on Ramanujan's continued fractions have been discussed.

Keywords and Phrases: Continued fraction, q-series, componendo and dividendo rule, q-binomial theorem.

2020 Mathematics Subject Classification: 33D15, 33D65.

1. Introduction, Notations and Definitions

The subject of continued fractions (CF) is an old subject although many people are not aware of it. Actually, continued fractions have so many applications in algebra and in various fields such as mathematics, physics, and chemistry. The easiest way of forming a continued fraction is by writing a certain amount in the form of a numerator and a denominator, and each denominator is composed of a numerator and a denominator and so on. Usually, the successive numerators are equal to one.

Continued fractions have a long history; they were known since the appearance of Euclidean algorithm for finding the greatest common divisor (GCD) of two numbers. That was around the year 300 B.C.. Research works and papers continue then to be performed and a huge accumulation of applications arise; this is due to their simplicity to deal with and the smooth way of the calculations involved. We should add that the subject of continued fractions is still very fruitful and interesting for researchers [4, 5, 6, 9, 10] all over the globe.