

**ADRIATIC INDICES OF SOME DERIVED GRAPHS OF
TRIGLYCERIDE**

V. Lokesha, M. Manjunath*, T. Deepika[¶], A. S. Cevik[†], I. N. Cangul[§]

Department of Studies in Mathematics,
Vijayanagara Sri Krishnadevaraya University,
Ballari - 583105, Karnataka, INDIA

E-mail : v.lokesha@gmail.com

*Department of Mathematics,
Ballari Institute of Technology and Management,
Ballari, Karnataka - 583105, INDIA

E-mail : manju3479@gmail.com

[¶]Department of Mathematics, School of Engineering,
Dayananda Sagar University, Bangalore, Karnataka - 560068, INDIA

E-mail : sastry.deepi@gmail.com

[†]Department of Mathematics,
Faculty of Science, Selcuk University 42075 Konya, TURKEY

E-mail : sinan.cevik@selcuk.edu.tr

[§]Department of Mathematics,
Bursa Uludag University 16059 Bursa, TURKEY

E-mail : cangul@uludag.edu.tr

(Received: Jan. 24, 2021 Accepted: Oct. 16, 2021 Published: Dec. 30, 2021)

Abstract: Topological indices are graph invariants used for quantitative structure-activity relationship (QSAR) and quantitative structure-property relationship (QSPR) in molecular chemistry. Esters are a class of organic compounds which are mostly fragrant and can be represented by the formula RCOOR. They are usually