

BILATERAL GENERATING RELATIONS FOR CERTAIN HYPERGEOMETRIC FUNCTIONS OF MORE THAN ONE VARIABLE

P.K.Banerji and F.B.F. Mohsen

Department of Mathematics, Faculty of Science
J.N.V. University, Jodhpur-342005
E-main : banerjipk@yahoo.com

(Received on November, 25, 2002)

Abstract : This paper establishes some bilateral generating relations for hypergeometric functions of two, three and four variables. Some particular cases have been explored.

1. Introduction. The hypergeometric functions of four variables namely K_i ($i = 9, 10, 11, 12, 13$) were studied by Exton [2, p.78], while a study of hypergeometric functions of three variables can be found in Srivastava and Manocha [6, pp. 66-68]. Horn's functions, can be found in Erdélyi et al. [1, p.225]. In what follows, we mention following results, those are required in our investigation.

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_1[-n, \mu, \nu; \alpha; x, y] t^n = (1-t)^{-\lambda} F_1\left[\lambda, \mu, \nu; \alpha; \frac{xt}{t-1}, \frac{yt}{t-1}\right], \quad (1.1)$$

where $\max\left\{\left|\frac{xt}{t-1}\right|, \left|\frac{yt}{t-1}\right|, |t|\right\} < 1$.

[cf. Srivastava [5]].

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_2[-n, \mu, \nu; \alpha, \beta; x, y] t^n = (1-t)^{-\lambda} F_2\left[\lambda, \mu, \nu; \alpha, \beta; \frac{xt}{t-1}, \frac{yt}{t-1}\right], \quad (1.2)$$

where $\max \left\{ \left| \frac{xt}{t-1} \right| + \left| \frac{yt}{t-1} \right|, |t| \right\} < 1$.

[cf. Srivastava [5]].

$$\begin{aligned} \sum_{n=0}^{\infty} \binom{v+n}{n} \frac{(\lambda)_n}{(\mu)_n} P_{n+v}^{(\alpha, \beta-n-v)}(x) t^n \\ = \binom{\alpha+v}{v} \left(\frac{1+x}{2} \right)^{v-\beta} F_2 \left[\alpha+v+1, -\beta, \lambda; \alpha+1, \mu; \frac{1-x}{2}, \frac{(x+1)t}{2} \right]. \end{aligned} \quad (1.3)$$

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{(\mu)_n} P_n^{(\alpha, \beta-n)}(x) t^n = F_1 \left[\lambda, -\beta, \lambda; \alpha+\beta+1, \mu; t, \frac{(x+1)t}{2} \right]. \quad (1.4)$$

Above two results are due to Srivastava [4].

Following four results are found in Srivastava and Manocha [6] :

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} {}_2F_1[\rho-n, \alpha; \beta; x] t^n = (1-t)^{-\lambda} F_1 \left[\alpha, \rho, \lambda; \beta; x, -\frac{xt}{1-t} \right]. \quad (1.5)$$

$$\sum_{n=0}^{\infty} \frac{(\beta)_n (\gamma)_n}{(\delta)_n} {}_2F_1[-n, \alpha; \beta; x] \frac{t^n}{n!} = F_1 \left[\gamma, \beta-\alpha, \alpha; \delta; t, (1-x)t \right]. \quad (1.6)$$

$$\begin{aligned} \sum_{n=0}^{\infty} \binom{m+n}{n} \frac{(\lambda)_n}{(-\alpha-\beta-m)_n} P_{m+n}^{(\alpha-n, \beta-n)}(x) t^n \\ = \binom{\alpha+\beta+2m}{m} \left(\frac{x+1}{2} \right)^m \left[1 + \frac{(x+1)t}{2} \right]^{-\lambda} \\ \times F_1 \left[-\beta-m, -m, \lambda; -\alpha-\beta-2m; \frac{2}{x+1}, \frac{t}{1+(x+1)t/2} \right] \end{aligned} \quad (1.7)$$

and

$$\sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} {}_2F_1[-n, a; b; x] t^n = (1-t)^{-\lambda} {}_2F_1 \left[\lambda, a; b; -\frac{xt}{1-t} \right]. \quad (1.8)$$

In above results, whenever they appear, F_1 and F_2 are Appell's functions and ${}_2F_1(\cdot)$ is the Gauss hypergeometric function.

2. Main Results and Proofs

We have established the following bilateral generating relations:

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_E [\lambda + n, \lambda + n, \lambda + n, \alpha, \beta, \beta; \gamma_1, \gamma_2, \gamma_3; x, y, z] F_1(-n, \mu, \nu; \gamma; u, v) t^n \\ = (1-t)^{-\lambda} \sum_{p=0}^{\infty} \frac{(\lambda)_p (\gamma)_p}{(\gamma_1)_p p!} \left(\frac{x}{1-t} \right)^p \\ \times K_9 \left[\lambda + p, \lambda + p, \lambda + p, \lambda + p, \beta, \beta, \mu, \nu; \gamma_2, \gamma_3, \gamma, \gamma; \frac{y}{1-t}, \frac{z}{1-t}, \frac{ut}{t-1}, \frac{vt}{t-1} \right]. \end{aligned} \quad (2.1)$$

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_F [\lambda + n, \lambda + n, \lambda + n, \alpha_1, \alpha_2, \alpha_1; \beta_1, \beta_2, \beta_2; x, y, z] F_2(-n, \mu, \nu; \alpha, \beta; u, v) t^n \\ = (1-t)^{-\lambda} \sum_{q=0}^{\infty} \frac{(\lambda)_q (\alpha_2)_q}{(\beta_2)_q q!} \left(\frac{y}{1-t} \right)^q \\ \times K_{10} \left[\lambda + q, \lambda + q, \lambda + q, \lambda + q, \alpha_1, \alpha_1, \mu, \nu; \beta_1, \beta_2 + q, \alpha, \beta; \frac{x}{1-t}, \frac{z}{1-t}, \frac{ut}{t-1}, \frac{vt}{t-1} \right]. \end{aligned} \quad (2.2)$$

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_D^{(3)} (\lambda + n, \alpha_1, \alpha_2, \alpha_3; \gamma, x, y, z) {}_2F_1(-n, a; b; u) t^n \\ = (1-t)^{-\lambda} K_{11} \left[\lambda, \lambda, \lambda, \lambda, \alpha_1, \alpha_2, \alpha_3, a; \gamma, \gamma, \gamma, b; \frac{x}{1-t}, \frac{z}{1-t}, \frac{ut}{t-1} \right]. \end{aligned} \quad (2.3)$$

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_G [\lambda + n, \lambda + n, \lambda + n, \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_2; x, y, z] P_n^{(\alpha, \beta-n)}(u) t^n \\ = \sum_{p=0}^{\infty} \frac{(\lambda)_p (\beta_1)_p}{(\gamma_1)_p p!} x^p \\ \times K_{12} [\lambda + p, \lambda + p, \lambda + p, \lambda + p, \beta_2, \beta_2 - \beta, \alpha + \beta + 1; \gamma_2, \gamma_2, \mu, \mu; y, z, t, 1/2(u+1)] . \end{aligned}$$

(2.4)

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_G[\lambda+n, \lambda+n, \lambda+n, \beta_1, \beta_2, \beta_3; \gamma_1, \gamma_2, \gamma_2; x, y, z] F_2[-n, \mu, \nu; \alpha, \beta; u, v] t^n \\
&= (1-t)^{-\lambda} \sum_{p=0}^{\infty} \frac{(\lambda)_p (\beta_1)_p}{(\gamma_1)_p p!} \left(\frac{x}{1-t} \right)^p \\
&\times K_{13} \left[\lambda+p, \lambda+p, \lambda+p, \lambda+p, \beta_2, \beta_3, \mu, \nu; \gamma_2, \gamma_2, \alpha, \beta; \frac{y}{1-t}, \frac{z}{1-t}, \frac{ut}{t-1}, \frac{vt}{t-1} \right].
\end{aligned}
\tag{2.5}$$

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{(\beta)_n (\gamma)_n}{(\delta)_n n!} H_3[\alpha', \beta'; \delta+n; x, y] {}_2F_1[-n, \alpha; \beta; u] t^n \\
&= \sum_{p=0}^{\infty} \frac{(\alpha')_{2p} x^p}{(\delta)_p p!} \\
&\times F_S[\alpha' + 2p, \gamma, \gamma, \beta', \beta - \alpha, \alpha; \delta + p, \delta + p, \delta + p; y, t, (1-u)t] \quad .
\end{aligned}
\tag{2.6}$$

$$\begin{aligned}
& \sum_{n=0}^{\infty} \binom{m+n}{n} \frac{(\lambda)_n}{(-\alpha-\beta-m)_n} H_3[\lambda+n; \gamma, \delta; x, y] P_{m+n}^{(\alpha-n, \beta-n)}(u) t^n \\
&= \binom{\alpha+\beta+2m}{m} \left(\frac{u-1}{2} \right)^m \left[1 + \frac{(u+1)t}{2} \right]^{-\lambda} \sum_{p=0}^{\infty} \frac{(\lambda)_{2p}}{(\delta)_p p!} \left[\frac{x}{\left\{ 1 + \frac{u+1}{2} \right\}^2} \right]^p \\
&\times F_M \left[\gamma, -\beta-m, -\beta-m, \lambda+2p, -m, \lambda+2p; \delta+p, -\alpha-\beta-2m, -\alpha-\beta-2m; \right. \\
&\quad \left. \frac{y}{1 + \frac{(u+1)t}{2}}, \frac{u}{u+1}, \frac{t}{1 + \frac{(u+1)t}{2}} \right] \quad .
\end{aligned}
\tag{2.7}$$

$$\begin{aligned}
& \sum_{n=0}^{\infty} \binom{v+n}{n} \frac{(\lambda)_n}{n!} H_4[\lambda+n; \beta'; \gamma, \delta; x, y] P_{n+v}^{(\alpha, \beta-n-v)}(u) t^n \\
&= \binom{\alpha+v}{v} \left(\frac{1+u}{2} \right)^{v-\beta} \sum_{p=0}^{\infty} \frac{(\lambda)_{2p} x^p}{(\gamma)_p p!} \\
&\times F_K \left[\beta', \alpha+v+1, \alpha+v+1, \lambda+2p, -\beta, \lambda+2p; \delta, \alpha+1, \mu; y, \frac{1-u}{2}, \frac{(1+u)t}{2} \right].
\end{aligned} \tag{2.8}$$

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} G_2[\lambda+n, \alpha', \beta, \beta'; x, y] {}_2F_1(p-n, \alpha; \gamma; u) t^n \\
&= (1-t)^{-\lambda} \sum_{q=0}^{\infty} \frac{(\alpha')_q (\beta)_q (-y)^q}{(1-\beta'-p)_q q!} F_M \left[\beta', \alpha, \alpha, \lambda, \rho, \lambda, 1-\beta-q, \gamma, \gamma, \frac{x}{t-1}, u, \frac{ut}{t-1} \right].
\end{aligned} \tag{2.9}$$

and

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} H_4[\lambda+n, \beta; \gamma, -\delta; x, y] F_2(-n, \mu, v; \alpha, \beta; u, v) t^n \\
&= (1-t)^{-\lambda} \sum_{q=0}^{\infty} \frac{(\lambda)_q (\beta)_q}{(\delta)_q q!} \left(\frac{y}{1-t} \right)^q X_8 \left[\lambda+q, \mu, v; \gamma, \alpha, \beta; \frac{x}{(1-t)^2}, \frac{ut}{t-1}, \frac{vt}{t-1} \right].
\end{aligned} \tag{2.10}$$

Proofs of the Results:

In the proofs of the above mentioned ten generating relations, we will use the relation

$$(\lambda)_n (\lambda+n)_{p+q+r} = (\lambda)_{n+p+q+r} = (\lambda)_{p+q+r} (\lambda+p+q+r)_n \tag{2.11}$$

Proof of (2.1):

Let us write A for the terms on the LHS of (2.1). Using (1.1) and (2.11), we have

$$A = \sum_{p,q,r=0}^{\infty} \frac{(\lambda)_{p+q+r} (\alpha)_p (\beta)_{q+r} x^p y^q z^r}{(\gamma_1)_p (\gamma_2)_q (\gamma_3)_r p! q! r!} (1-t)^{-(\lambda+p+q+r)} F_1 \left(\lambda + p + q + r, \mu, \nu, r; \frac{ut}{t-1}, \frac{vt}{t-1} \right). \quad (2.12)$$

Now expressing $F_1(\cdot)$ in series and using (2.11), we get

$$A = (1-t)^{-\lambda} \sum_{p=0}^{\infty} \frac{(\lambda)_p (r)_p}{(r_1)_p p!} \left(\frac{x}{1-t} \right)^p \sum_{q,r,i,j=0}^{\infty} \frac{(\lambda+p)_{q+r+i+j} (\beta)_{q+r}}{(r_2)_q (r_3)_r (r)_{i+j}} \times \frac{(\mu)_i (\nu)_j}{q! r! i! j!} \left(\frac{y}{1-t} \right)^q \left(\frac{z}{1-t} \right)^r \left(\frac{ut}{t-1} \right)^i \left(\frac{vt}{t-1} \right)^j. \quad (2.13)$$

By virtue of the expression for K_9 [cf. Exton [2, p.78]], the equation (2.13) yields the required result.

Employing (1.2), (1.4) and (1.8) the relations (2.2) - (2.5) will be obtained following above procedure.

Proof of (2.6):

Replace by B the terms on the LHS of (2.6) and then using (1.6) and (2.11), we get

$$B = \sum_{p,q=0}^{\infty} \frac{(\alpha')_{2p+q} (\beta')_q}{(\delta)_{p+q} p! q!} x^p y^q F_1(\gamma, \beta - \alpha, \alpha; \delta + p + q; t, (1-u)t) .$$

Now write F_1 in terms of series and employ (2.11), we have

$$B = \sum_{p=0}^{\infty} \frac{(\alpha')_{2p}}{(\delta)_p p!} x^p \sum_{q,i,j=0}^{\infty} \frac{(\alpha' + 2p)_q (r)_{i+j} (\beta')_q (\beta - \alpha)_i (\alpha)_j}{(\delta + p)_{q+i+j} q! i! j!} y^q t^i [(1-ut)^j] .$$

(2.14)

The result is obtained by using [6, p.68].

Results (2.7) – (2.10) are obtained in the manner (2.6) is obtained, we use (1.7), (1.3), (1.5) and (1.2) respectively.

3. Particular Cases

The particular cases, believed to be interesting, give bilateral generating relations for Appell's functions, for Laguerre polynomial and for the confluent form of F_2 , ψ_1 and ψ_2 [see e.g. [6], p. 59].

(i) In (2.1), put $x = 0$, we then get

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_4[\lambda + n, \beta, \gamma_2, \gamma_3; y, z] F_1(-n, \mu, \nu; \gamma, u, v) t^n \\ &= (1-t)^{-\lambda} K_9 \left[\lambda, \lambda, \lambda, \lambda, \beta, \mu, \nu, \gamma_2, \gamma_3, \gamma, \gamma, \frac{y}{1-t}, \frac{z}{1-t}, \frac{ut}{t-1}, \frac{vt}{t-1} \right]. \end{aligned} \quad (3.1)$$

(ii) Now, if in (2.1), $\nu = 0, \gamma_3 \rightarrow 0$ and if y is replaced by y/β and u by u/μ , then taking limit as $\mu, \beta \rightarrow \infty$, we have

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} \psi_1[\lambda + n, \alpha; \gamma_1, \gamma_2; x, y] {}_1F_1(-n; \gamma; u) t^n \\ &= (1-t)^{-\lambda} \sum_{p=0}^{\infty} \frac{(\lambda)_p (\gamma)_p}{(\gamma_1)_p p!} \left(\frac{x}{1-t} \right)^p \psi_2 \left[\lambda + p, \gamma_1, \gamma_2, \frac{ut}{t-1}, \frac{y}{1-t} \right] \end{aligned} \quad (3.2)$$

(iii) For $\beta_1 = 0$, (2.4) yields

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(\lambda)_n}{n!} F_1[\lambda + n, \beta_2, \beta_3, \gamma_2; y, z] P_n^{(\alpha, \beta-n)}(u) t^n \\ &= K_{12}[\lambda, \lambda, \lambda, \lambda, \beta_2, \beta_2, -\beta, \alpha + \beta + 1, \gamma_2, \gamma_2, \mu, \mu; y, z, t, 1/2(u+1)t] \end{aligned} \quad (3.3)$$

(iv) In (2.5) if we put $\beta_1 = \beta_3 = \nu = 0$ and replace u by u/μ , by taking limit as $|\mu| \rightarrow \infty$ and putting $\alpha = \lambda$, we get

$$\sum_{n=0}^{\infty} {}_2F_1(\lambda+n, \beta_2; \gamma_2; y) L_n^{(\lambda-1)}(u) t^n = (1-t)^{-\lambda} \psi_1 \left[\lambda, \beta_2; \gamma_2, \lambda; \frac{y}{1-t}, \frac{ut}{t-1} \right]. \quad (3.4)$$

Acknowledgements

This work is partially supported by the Research Scheme of CSIR (India) No. 25(0105)/99/ EMR-II and Major Research Project of UGC (India) No. F.8-24/2001/SR-I sanctioned to PKB.

References

1. Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. Higher Transcendental Functions, Vol. I, McGraw – Hill Book Co. Inc., New York, London (1953).
2. Exton, H. Multiple Hypergeometric Functions and Applications, John – Wiley & Sons, New York (1976).
3. Exton, H. J. Indian Acad. Math, 4 (1982), 113-119.
4. Srivastava, H. M. On generating function for the Jacobi polynomials, J. Math. Sci. 4 (2) (1969), 61-68.
5. Srivastava, H. M. Certain formulas involving Appell's functions, Comment. Math. Univ. St. Pauli, 21 (1) (1972), 73-99.
6. Srivastava, H. M. and Manocha, H. L. A Treatise on Generating Functions, John-Wiley and Sons, New York (1984).